



**PCR BIOSYSTEMS**  
simplifying research

## IsoFast™ Bst Polymerase (120 U/μL)

[www.pcrbio.com](http://www.pcrbio.com)

### Product description

IsoFast™ Bst Polymerase is a recombinant protein expressed in *E. coli* and represents the large fragment of *Geobacillus stearothermophilus* (formerly known as *Bacillus stearothermophilus*) DNA Polymerase. This portion of the protein maintains 5' to 3' polymerase activity but lacks the 5' to 3' exonuclease activity.<sup>1</sup>

IsoFast™ Bst Polymerase displays strong strand displacement activity and is suitable for nucleic acid amplification methods such as whole genome amplification, multiple displacement amplification and isothermal amplification. We recommend a reaction temperature of 65 °C. However, the enzyme works well over a broad temperature range, from 55 °C to 70 °C. It can be heat inactivated at 80 °C.

Designed for fast amplification speed, IsoFast™ Bst Polymerase gives rapid and consistent results across different target sequences and sample types. The enzyme is provided with a 2-part buffer system to ensure high yield and performance even under difficult conditions.

Real-time detection with any qPCR thermocycler can be achieved by adding 20x Fluorescent Dye to the reaction. This is supplied separately, catalogue number PB80.30-2 and PB80.30-10.

The high concentration format of this enzyme makes it ideal for applications requiring little buffer carry over (lyophilisation or air-drying) and for highthroughput screening workflows.

Component	8000 units	150,000 units	1,500,000 units
IsoFast Bst Polymerase (120 U/μL)	1 x 67 μL	1 x 1.25 mL	1 x 12.5 mL
10x IsoFast Buffer A	2 x 1.25 mL	1 x 47 mL	1 x 470 mL
5x IsoFast Buffer B	3 x 1.7 mL	1 x 94 mL	1 x 940 mL

### Shipping and storage

On arrival the kit should be stored between -30 °C and -15 °C. Keep components on ice when in use. If stored correctly the kit will retain full activity for 12 months. We recommend aliquoting the enzyme at the first use to avoid excess freeze/thaw cycles.

### Limitations of product use

The product may be used for in vitro research purposes only.

### Technical support

Help and support is available on our website at <https://pcrbio.com/resources/> including answers to frequently asked technical questions. For technical support and troubleshooting you can submit a technical enquiry online, or alternatively email [technical@pcrbio.com](mailto:technical@pcrbio.com) with the following information:

- Amplicon size
- Reaction setup
- Reaction conditions
- Screen grabs or images of amplification results

1. Mead DA, McClary JA, Luckey JA, Kostichka AJ, Witney FR, Smith LM. Bst DNA polymerase permits rapid sequence analysis from nanogram amounts of template. *Biotechniques*. 1991 Jul;11(1):76-8, 80, 82-87.

## Important considerations

**10x IsoFast Buffer A:** The 10x buffer contains 30 mM MgSO<sub>4</sub>, 16 mM dNTPs, enhancers and stabilizers. The buffer composition has been optimised to maximise the rate of amplification.

**5x IsoFast Buffer B:** The 5x buffer contains enhancers designed to further increase the reaction speed.

## Example usage: Strand displacement

Reaction temperature	Reaction time	Deactivation temperature	Deactivation time
Recommended: 65 °C Optimal range: 55-70 °C	30-60 minutes	80 °C	10 minutes

## Example usage: Loop-mediated isothermal amplification (LAMP)

1. Allow each component to reach room temperature, then briefly vortex.
2. Prepare a master mix based on the following table. Reactions should be set up on ice:

Reagent	25 µL reaction	Final concentration	Notes
10x IsoFast Buffer A	2.50 µL	1x	
5x IsoFast Buffer B	5.00 µL	1x	
20x Fluorescent Dye (optional)	1.25 µL	1x	Available separately, catalogue number PB80.30-02 and PB80.30-10
IsoFast Bst Polymerase (120 U/µL)	0.067 µL	8 U	
10x Primer set	2.50 µL	1x	We recommend a predicted melting temperature of around 60 °C using default Primer Explorer v5 settings. A primer set can be prepared with all 4 or 6 (if you include Loop) primers. A 10x primer set should contain: 16 µM FIP, 16 µM BIP, 2 µM F3, 2 µM B3, 4-8 µM LoopF, 4-8 µM LoopB in TE Buffer or water.
Template DNA	Variable		
PCR grade dH <sub>2</sub> O	Up to 25 µL final volume		

3. Incubate at 65 °C for 30 minutes. Time can be extended and temperature can be modified (between 55 °C and 70 °C) as necessary for low copy targets, challenging templates, or whenever amplification times have been reported to be slow.

If a qPCR instrument is used for signal detection, follow the reaction using the FAM channel, acquiring data every 10-15 seconds. If final products are to be analysed after the reaction is complete, the enzyme can be inactivated by heating at 80 °C for 10 minutes.

## Notice to purchasers

Some uses of this product may be the subject of patent rights. No licence or rights are conveyed to the purchaser of this product. It is the sole responsibility of the buyer to ensure that use of the product does not infringe any patent rights of third parties.